

New Approach to Texture-zeros with S_3 symmetry

- Flavor Symmetry and Vacuum Aligned Mass Textures - *

Satoru Kaneko^{1,†}, Hideyuki Sawanaka^{2,‡}, Takaya Shingai^{2,§},
Morimitsu Tanimoto^{3,¶} and Koichi Yoshioka^{4,||}

¹*AHEP Group, Institut de Física Corpuscular–C.S.I.C/Universitat de València,
Edifici Instituts d'Investigació, Apt. 22085, E-46071 València, Spain*

²*Graduate school of Science and Technology, Niigata University, Niigata 950-2181, Japan*

³*Physics Department, Niigata University, Niigata 950-2181, Japan*

⁴*Physics Department, Kyushu University, Fukuoka 812-8581, Japan*

Abstract

A texture-zeros is an approach to reduce the number of free parameters in Yukawa couplings and it is one of the most attractive ones. In our paper, we discuss the origin of zero-structure in texture-zeros by S_3 flavor symmetry approach. Some of electroweak doublet Higgs fields have vanishing vacuum expectation value (VEV) which leads to vanishing elements in quark and lepton mass matrices. Then, the structure of supersymmetric scalar potential is analyzed and Higgs fields have non-trivial S_3 charges. As a prediction of our paper, a lower bound of a MNS matrix element, $U_{e3} \geq 0.04$, is obtained. The suppression of flavor-changing neutral currents (FCNC) mediated by the Higgs fields is discussed and lower bounds of the Higgs masses are derived.

1 Introduction

It is known that the standard model of elementary particle physics has many free Yukawa couplings. The texture-zeros was considered as an approach to reduce the number of the Yukawa couplings. By this approach, predictions among observable values are obtained. Due to useful features of the texture-zeros, many authors have studied the texture-zeros [1]–[10]. However, an origin of zero elements of mass matrices are not known. Namely, such zeros have often been assumed by hand. Our paper proposes a new approach to solve this problem and some models are considered by $U(1)$ flavor symmetry [11] or discrete flavor symmetry [12].

In our model, we take the discrete flavor symmetry approach. Flavor symmetry is expected to be a clue to understand the masses and mixing angles of quarks and leptons. It reduces the number of free parameters in Yukawa couplings such as the texture-zeros and some testable predictions of masses and mixing angles generally follow. Some predictive models with discrete flavor symmetries have been explored by many authors [13]–[20]. We consider a S_3 group as the discrete symmetry because S_3 group is the simplest discrete non-Abelian group. S_3 is a permutation group of three objects.

In this paper, we discuss the mass matrix of quarks and leptons in the case that $SU(2)$ doublet Higgs fields have non-trivial S_3 flavor charges. Then, we discuss a dynamical realization of the texture-zeros. In previous models, some Yukawa couplings are restricted by discrete

*talked by T. Shingai at the International Workshop on Neutrino Masses and Mixings — Toward Unified Understanding of Quark and Lepton Mass Matrices —, University of Shizuoka, Shizuoka, Japan, December 17–19 2006.

[†]satoru@ific.uv.es

[‡]hide@muse.sc.niigata-u.ac.jp

[§]shingai@muse.sc.niigata-u.ac.jp

[¶]tanimoto@muse.sc.niigata-u.ac.jp

^{||}yoshioka@higgs.phys.kyushu-u.ac.jp

flavor symmetry and then some favored textures(texture-zeros) are obtained by changing basis. However, there is another possibility to obtain zero matrix elements. This is vanishing Higgs VEV. This possibility, however, does not make sense if a model contains only one Higgs field. Therefore, we introduce some Higgs fields. We perform the analysis of Higgs potential at the electroweak scale and examine whether some of Higgs fields in the flavor basis have vanishing expectation values. Vanishing elements(texture-zeros) of fermion mass matrices are obtained dynamically in the vacuum of the theory, and their positions are controlled by the flavor symmetry. An interesting point of the scheme is that, due to the S_3 group structure, the up and down quark mass matrices automatically have different forms at the vacuum, which lead to non-vanishing generation mixing.

It is found that the exact S_3 model has Nambu-Goldstone bosons. However, these disappear by introducing S_3 soft breaking terms and it is showed that these terms have no effect on predictions of the exact S_3 model. Generally speaking, S_3 soft breaking terms give rise to changing conditions necessary to take a favored vacuum. However, S_3 soft breaking terms introduced in our model do not generate conditions for a favored vacuum. As a prediction for a unitary matrix element U_{e3} , we obtain $U_{e3} \geq 0.04$. This will be tested in future experiments such as the double Chooz. The suppression of FCNC mediated by multiple Higgs fields will also be discussed.

2 S_3 invariant mass matrix on Supersymmetry

In this section, the S_3 invariant mass matrices are presented[14]. We consider a supersymmetric theory and both left-handed and right-handed fermions transform under a single S_3 symmetry. S_3 has three irreducible representations, a singlet, a pseudo singlet and a doublet. We suppose that two of three generations belong to S_3 doublets and the others are singlets. Namely, S_3 doublet is assigned to matter superfield of first and second generation, $\Psi_{L_{1,2}}$ and $\Psi_{R_{1,2}}$. Then, the most general supersymmetric Yukawa mass terms are obtained as

$$W_y = \Psi_{Li}(M_D)_{ij}\Psi_{Rj}, \quad (1)$$

$$M_D = \left(\begin{array}{cc|c} aH_1 & bH_S + cH_A & dH_2 \\ bH_S - cH_A & aH_2 & dH_1 \\ \hline eH_2 & eH_1 & fH_S \end{array} \right), \quad (2)$$

where $a, b, \dots, f$ are independent Yukawa coupling constants. H_1 and H_2 are S_3 doublet Higgs, H_A is S_3 pseudo singlet Higgs and H_S is S_3 singlet Higgs. Eq. (2) is obtained by using a decomposition of a tensor product of S_3 doublet, ϕ and ψ , which is given as

$$\phi^c \times \psi = (\phi_2\psi_2, \phi_1\psi_1)_2^T + (\phi_1\psi_2 - \phi_2\psi_1)_{1_A} + (\phi_1\psi_2 + \phi_2\psi_1)_{1_S}, \quad (3)$$

where $\phi^c = \sigma_1\phi^* = \sigma_1(\phi_1^*, \phi_2^*)^T$, $\psi = (\psi_1, \psi_2)^T$. Now we assign 1_S to third generation temporarily. We reconfigure this assignment later.

The S_3 invariant bare Majorana mass matrix for matter superfield $\Psi_i(i = 1, 2, 3)$ is given as

$$W_m = \Psi_i(M_R)_{ij}\Psi_j, \quad (4)$$

$$M_R = \left(\begin{array}{c|c} M_1 & \\ \hline M_1 & \\ \hline & M_2 \end{array} \right), \quad (5)$$

where M_1, M_2 are majorana masses.

3 S_3 invariant Higgs scalar potential analysis

In our model we consider eight Higgs bosons: H_{uS} , H_{dS} , H_{uA} , H_{dA} , H_{u1} , H_{d1} , H_{u2} and H_{d2} , which are $SU(2)$ weak doublet. Our purpose in this section is to discuss whether or not there are some vanishing VEV patterns with no parameter relations. S_3 invariant potential is given as

$$V = V_{\text{susy}} + V_{\text{soft}} , \quad (6)$$

$$\begin{aligned} V_{\text{susy}} = & \frac{1}{2}D_Y^2 + \frac{1}{2} \sum_{a=1,2,3} (D^a)^2 \\ & + |\mu_S|^2 (H_{uS}^\dagger H_{uS} + H_{dS}^\dagger H_{dS}) + |\mu_A|^2 (H_{uA}^\dagger H_{uA} + H_{dA}^\dagger H_{dA}) \\ & + |\mu_D|^2 (H_{u1}^\dagger H_{u1} + H_{d1}^\dagger H_{d1} + H_{u2}^\dagger H_{u2} + H_{d2}^\dagger H_{d2}) , \end{aligned} \quad (7)$$

$$\begin{aligned} V_{\text{soft}} = & m_{uS}^2 H_{uS}^\dagger H_{uS} + m_{dS}^2 H_{dS}^\dagger H_{dS} + (b_S H_{uS} \epsilon H_{dS} + \text{h.c.}) \\ & + m_{uA}^2 H_{uA}^\dagger H_{uA} + m_{dA}^2 H_{dA}^\dagger H_{dA} + (b_A H_{uA} \epsilon H_{dA} + \text{h.c.}) \\ & + m_{uD}^2 (H_{u1}^\dagger H_{u1} + H_{u2}^\dagger H_{u2}) + m_{dD}^2 (H_{d1}^\dagger H_{d1} + H_{d2}^\dagger H_{d2}) \\ & + [b_D (H_{u1} \epsilon H_{d2} + H_{u2} \epsilon H_{d1}) + \text{h.c.}] , \end{aligned} \quad (8)$$

where ϵ is the antisymmetric tensor, and b_x , m_{ux} and m_{dx} ($x = S, A, D$) are the holomorphic and non-holomorphic mass parameters of supersymmetry breaking, respectively. The D-terms are explicitly given by

$$\begin{aligned} D_Y = & -\frac{1}{2}g_Y \left\{ (H_{uS}^\dagger H_{uS} - H_{dS}^\dagger H_{dS}) + (H_{uA}^\dagger H_{uA} - H_{dA}^\dagger H_{dA}) \right. \\ & \left. + (H_{u1}^\dagger H_{u1} - H_{d1}^\dagger H_{d1}) + (H_{u2}^\dagger H_{u2} - H_{d2}^\dagger H_{d2}) \right\} , \end{aligned} \quad (9)$$

$$\begin{aligned} D^a = & -g_2 \left\{ (H_{uS}^\dagger T^a H_{uS} + H_{dS}^\dagger T^a H_{dS}) + (H_{uA}^\dagger T^a H_{uA} + H_{dA}^\dagger T^a H_{dA}) \right. \\ & \left. + (H_{u1}^\dagger T^a H_{u1} + H_{d1}^\dagger T^a H_{d1}) + (H_{u2}^\dagger T^a H_{u2} + H_{d2}^\dagger T^a H_{d2}) \right\} , \end{aligned} \quad (10)$$

with T^a ($a = 1, 2, 3$) is the $SU(2)_L$ generators. Hence, we analyze eight equations at vacuum:

$$(|\mu_S|^2 + m_{uS}^2)v_{uS} = b_S v_{dS} - X v_{uS}, (|\mu_S|^2 + m_{dS}^2)v_{dS} = b_S v_{uS} + X v_{dS}, \quad (11)$$

$$(|\mu_A|^2 + m_{uA}^2)v_{uA} = b_A v_{dA} - X v_{uA}, (|\mu_A|^2 + m_{dA}^2)v_{dA} = b_A v_{uA} + X v_{dA}, \quad (12)$$

$$(|\mu_D|^2 + m_{uD}^2)v_{u1} = b_D v_{d2} - X v_{u1}, (|\mu_D|^2 + m_{dD}^2)v_{d2} = b_D v_{u1} + X v_{d2}, \quad (13)$$

$$(|\mu_D|^2 + m_{uD}^2)v_{u2} = b_D v_{d1} - X v_{u2}, (|\mu_D|^2 + m_{dD}^2)v_{d1} = b_D v_{u2} + X v_{d1}, \quad (14)$$

where the quantities v_x are the absolute values of the neutral Higgs scalars,

$$\begin{aligned} v_{uS} &= |\langle H_{uS}^0 \rangle|, \quad v_{uA} = |\langle H_{uA}^0 \rangle|, \quad v_{u1} = |\langle H_{u1}^0 \rangle|, \quad v_{u2} = |\langle H_{u2}^0 \rangle|, \\ v_{dS} &= |\langle H_{dS}^0 \rangle|, \quad v_{dA} = |\langle H_{dA}^0 \rangle|, \quad v_{d1} = |\langle H_{d1}^0 \rangle|, \quad v_{d2} = |\langle H_{d2}^0 \rangle|, \end{aligned} \quad (15)$$

and the parameters b_S , b_A and b_D have been chosen to be real and positive using field redefinitions. X is defined as

$$X \equiv \frac{g_Y^2 + g_2^2}{4} (v_{uS}^2 - v_{dS}^2 + v_{uA}^2 - v_{dA}^2 + v_{u1}^2 - v_{d1}^2 + v_{u2}^2 - v_{d2}^2), \quad (16)$$

where g_Y and g_2 are gauge couplings of $U(1)_Y$ and $SU(2)_L$, respectively. In general, these equations are the coupled equations through a common parameter X which contains all the Higgs VEVs. However, we can separate these equations into three parts for the singlet, the pseudo singlet and the doublet. This is because vanishing-VEVs makes the equations trivial within each sector. Consequently, possible 14 VEV patterns in Table 1 are obtained. They do not need some parameter conditions.

v_{uS}	v_{dS}	v_{uA}	v_{dA}	v_{u1}	v_{u2}	v_{d1}	v_{d2}
0	0	0	0	0			0
0	0	0	0		0	0	
0	0	0	0				
0	0			0	0	0	0
0	0			0			0
0	0				0	0	
0	0						
		0	0	0	0	0	0
		0	0	0			0
		0	0		0	0	
		0	0				
				0	0	0	0
				0			0
					0	0	

Table 1: All possible minima of the scalar potential for S_3 singlet and doublet Higgs fields without tuning of Lagrangian parameters for electroweak symmetry breaking. The blank entries denote non-vanishing VEVs.

4 Quark and lepton mass textures

We obtained 14 VEV patterns in previous section. Now let us analyze these patterns phenomenologically. The most interesting pattern of 14 patterns is the following:

$$v_{uS} = v_{dS} = v_{u1} = v_{d2} = 0, \quad v_{uA}, v_{dA}, v_{u2}, v_{d1} \neq 0. \quad (17)$$

This pattern leads to the simplest texture(i.e.the maximal number of zero matrix elements) with non-trivial flavor mixing.

We consider mass matrices obtained from the VEV pattern (17). In section 2, we assigned S_3 singlet $\mathbf{1}_S$ to third generation. However, we can assign $\mathbf{1}_S$ to any generation in general. For example, $\mathbf{1}_S$ can be also assigned to first generation. As results of exhausting all the S_3 charge assignments, mass matrices are derived as

$$M_u = \begin{pmatrix} & b_u & \\ d_u & & f_u \\ & -f_u & i_u \end{pmatrix}, \quad M_d = \begin{pmatrix} & b_d & \\ d_d & e_d & \\ -e_d & & i_d \end{pmatrix}, \quad M_e = \begin{pmatrix} & d_d & 3e_d \\ b_d & -3e_d & \\ & & i_d \end{pmatrix}, \quad (18)$$

$$M_\nu = \begin{pmatrix} & b_\nu & c_\nu \\ -b_\nu & e_\nu & \\ g_\nu & & \end{pmatrix}, \quad M_R = \begin{pmatrix} & M_1 & \\ M_1 & & \\ & & M_2 \end{pmatrix}, \quad (19)$$

where blank entries denote zero and each parameter in M_u, M_d, M_e, M_ν such as d_u, d_d, b_ν stands for a product of a Yukawa coupling and a VEV, for example, $d_u = dv_{u2}$. We assume here the $SU(5)$ grand unification. Namely, S_3 charge assignment of (18) and (19) is embedded in the $SU(5)$ grand unification. It is, however, noted that the charge assignment of the right-handed neutrinos is completely irrelevant to low-energy physics, and the generation structure of the light neutrino mass matrix is determined only by the flavor charge of the left-handed leptons. We have included a group theoretical factor -3 [21] in front of the element e_d . Such a

Table 2: The $U(1)$ symmetries of the Higgs scalar potential with the S_3 breaking terms (21).

	H_{uS}	H_{dS}	H_{uA}	H_{dA}	H_{u1}	H_{u2}	H_{d1}	H_{d2}
U_X	+1	-1	0	0	+1	0	0	-1
U_Y	+1	-1	+1	-1	+1	+1	-1	-1

factor originates from a Yukawa coupling to higher-dimensional Higgs field. For example, this originates from 45-plet of $SU(5)$.

Some predictions from our model are given as

$$|V_{cb}| = \sqrt{\frac{m_c}{m_t}}, \quad |U_{e3}| \geq 0.04. \quad (20)$$

The former prediction in (20) is known to accurately fit the experimental data and it has already been studied in other theoretical framework [9, 22]. A precise numerical estimation, however, indicates that the former prediction is not satisfied with the experimental data slightly. There is a possibility to modify this deviation. That is a supersymmetric threshold correction. The theory would become viable in light of the current experimental data. We leave a detailed discussion of this correction to future investigations. The latter prediction[23] in (20) is most important prediction in our model. This lower bound will be tested in future experiments, such as the double Chooz[24].

5 Higgs mass spectrum and S_3 soft breaking terms in B-term

The S_3 potential has an enhanced global symmetry $SU(2) \times U(1)^2$ and leads to massless Nambu-Goldstone bosons in the electroweak broken phases. Global symmetries $U(1)^2$ is presented in Table 2. It is therefore reasonable to softly break the flavor symmetry within the scalar potential. We introduce the following supersymmetry-breaking soft terms which do not break phenomenological characters of the exact S_3 model.

$$V_{\mathcal{S}_3} = b_{SD} H_{uS} H_{d2} + b'_{SD} H_{u1} H_{dS} + b_{AD} H_{uA} H_{d1} + b'_{AD} H_{u2} H_{dA} + \text{h.c.} \quad (21)$$

A S_3 soft breaking model have the same phenomenological characters as the exact S_3 model and we can take the VEV pattern (17) with no parameter condition.

The S_3 soft breaking model also predicts a lightest Higgs mode. This is similar to the minimal supersymmetric standard model(MSSM). We obtain the neutral Higgs masses squared approximately:

$$\begin{aligned} & \{M_{h^0}^2, M_{H_1^0}^2, M_{H_2^0}^2, M_{H_3^0}^2, M_{H_4^0}^2\} \\ & = \{\mathcal{O}(v^2), \bar{m}^2 - \bar{b} - \bar{b}', \bar{m}^2 + \bar{b} - \bar{b}', \bar{m}^2 - \bar{b} + \bar{b}', \bar{m}^2 + \bar{b} + \bar{b}'\}. \end{aligned} \quad (22)$$

The other three mass eigenvalues squared take the complicated forms. Then, we make the simplifying assumption

$$|\mu_x|^2 + m_{ux}^2 = |\mu_x|^2 + m_{dx}^2 \equiv \bar{m}^2, \quad b_x \equiv \bar{b}, \quad (x = S, A, D) \quad (23)$$

$$b_{SD} = b'_{SD} = b_{AD} = b'_{AD} \equiv \bar{b}', \quad (24)$$




Figure 1: A Higgs-mediated tree-level FCNC process for the K meson system

with the hierarchy $\bar{m}^2, |\bar{b}|, |\bar{b}'| \gg v^2 \equiv v_{u2}^2 + v_{d1}^2 + v_{uA}^2 + v_{dA}^2$. Hence, we obtain a lightest mode which has a weak scale mass $\mathcal{O}(v)$ and its eigenvector is given as

$$h^0 = \frac{1}{v} \left(v_{uA} h_{uA}^0 + v_{dA} h_{dA}^0 + v_{u2} h_{u2}^0 + v_{d1} h_{d1}^0 \right). \quad (25)$$

6 $K^0 - \bar{K}^0$ mixing

Since there are multiple electroweak doublet Higgs bosons which couple to matter fields, flavor-changing processes are mediated at classical level by these Higgs fields. We can show that flavor-changing process at classical level leads to all but one Higgs masses which are order of supersymmetric breaking scale. Therefore, the experimental observations of FCNC rare events would show a bound on the supersymmetry breaking scale. Among various experimental constraints, we find the most important constraint comes from the neutral K meson mixing. For the heavy Higgs modes, the tree-level $K_L - K_S$ mass difference Δm_K^{tree} is given by the matrix element of the effective Hamiltonian between K mesons[25]. Δm_K^{tree} includes M_H , which is an average of the Higgs masses $1/M_H^2 = \frac{1}{4} \left(1/M_{H_1^0}^2 + 1/M_{H_2^0}^2 + 1/M_{H_3^0}^2 + 1/M_{H_4^0}^2 \right)$, and a free parameter η is given as

$$\eta = \frac{(y_d^S)_{22} b_d v_{d1}}{m_b^2} - \frac{(y_d^S)_{13} d_d v_{d1}}{m_s m_b}. \quad (26)$$

$(y_d^S)_{22}$ and $(y_d^S)_{13}$ are the down type quark Yukawa couplings. It is found from Fig. 2 that heavy Higgs masses are bounded from below so as to suppress the extra Higgs contribution compared with the standard model one. M_H is roughly given by

$$M_H \geq \begin{cases} 3.8 \text{ TeV} & (\eta = 0) \\ 1.4 \text{ TeV} & (\eta = 0.03) \end{cases}, \quad (27)$$

where we took $\eta = 0$ and 0.03 as typical values. We used the experimental values $m_K = 490 \text{ MeV}$ and $f_K = 160 \text{ MeV}$, and took $v_{d1} = 100 \text{ GeV}$ as a typical electroweak scale.

7 Summary

The mass matrix of quarks and leptons have discussed in view of S_3 flavor symmetry approach. We have analyzed the structure of the Higgs scalar potential and examined possible vanishing




Figure 2: $\Delta m_K^{\text{tree}} / \Delta m_K^{\text{SM}}$ as a function of the Higgs parameter M_H . The solid and dashed curves correspond to $\eta = 0$ and 0.03, respectively.

elements of quark and lepton mass matrices. Then, there exists the most favored VEV pattern for which parameter conditions are not necessary and quark and lepton mass matrices from the VEV pattern is consistent with experimental data. S_3 assignment to the flavor can be embedded in the $SU(5)$ grand unification and we take non-trivial S_3 assignment. As phenomenological results, a prediction for the lepton mixing U_{e3} is obtained: $U_{e3} \geq 0.04$. This prediction will be tested in future experiments. It is found from tree-level FCNC processes mediated by the Higgs bosons that TeV scale typical Higgs mass is consistent with K^0 - $\bar{K}^0$ mixing experimental bound.

Acknowledgements

We would like to thank the organizers of the International Workshop on Neutrino Masses and Mixings and thank J. Kubo, T. Kobayashi and H. Nakano for helpful discussions.

References

- [1] P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. **B548** (2002), 119.
M. Raidal and A. Strumia, Phys. Lett. **B553** (2003), 72.
Z. Xing, Phys. Rev. **D69** (2004), 013006.
G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim and T. Yanagida, Phys. Lett. **B562** (2003), 265.
- [2] H. Fritzsch, Phys. Lett. **B73** (1978), 317; Nucl. Phys. **B155** (1979), 189.
P. Ramond, R. G. Roberts and G. G. Ross, Nucl. Phys. **B406** (1993), 19.
- [3] H. Nishiura, K. Matsuda and T. Fukuyama, Phys. Rev. **D60** (1999), 013006.
E. K. Akhmedov, G. C. Branco and M. N. Rebelo, Phys. Rev. Lett. **84** (2000), 3535.
S. K. Kang and C. S. Kim, Phys. Rev. **D63** (2001), 113010.
- [4] P. H. Frampton, S. L. Glashow and D. Marfatia, Phys. Lett. **B536** (2002), 79.
Z.-z. Xing, Phys. Lett. **B530** (2002), 159.

- R. Barbieri, T. Hambye and A. Romanino, J. High Energy Phys. **03** (2003), 017.
A. Ibarra and G. G. Ross, Phys. Lett. **B575** (2003), 279.
- [5] A. Kageyama, S. Kaneko, N. Shimoyama and M. Tanimoto, Phys. Lett. **B538** (2002), 96.
S. Kaneko and M. Tanimoto, Phys. Lett. **B551** (2003), 127.
M. Honda, S. Kaneko and M. Tanimoto, J. High Energy Phys. **09** (2003), 028; Phys. Lett. **B593** (2004), 165.
S. Kaneko, H. Sawanaka and M. Tanimoto, J. High Energy Phys. **08** (2005), 073.
- [6] M.-C. Chen and K.T. Mahanthappa, Phys. Rev. **D68** (2003), 017301.
M. Bando and M. Obara, Prog. Theor. Phys. **109** (2003), 995.
M. Bando, S. Kaneko, M. Obara and M. Tanimoto, Phys. Lett. **B580** (2004), 229.
- [7] L. Lavoura, Phys. Lett. **B609** (2005), 317.
W. Grimus, S. Kaneko, L. Lavoura, H. Sawanaka and M. Tanimoto, J. High Energy Phys. **01** (2006), 110.
- [8] M. Frigerio and A. Y. Smirnov, Phys. Rev. **D67** (2003), 013007.
- [9] N. Uekusa, A. Watanabe and K. Yoshioka, Phys. Rev. **D71** (2005), 094024.
A. Watanabe and K. Yoshioka, J. High Energy Phys. **05** (2006), 044.
- [10] C. Hagedorn and W. Rodejohann, J. High Energy Phys. **07** (2005), 034.
A. Merle and W. Rodejohann, Phys. Rev. **D73** (2006), 073012.
- [11] L. Ibáñez and G. G. Ross, Phys. Lett. **B332** (1994), 100.
P. Binétruy, S. Lavignac and P. Ramond, Phys. Lett. **B350** (1995), 49; Nucl. Phys. **B477** (1996), 353.
P. Binétruy, S. Lavignac, S. T. Petcov and P. Ramond, Nucl. Phys. **B496** (1997), 3.
J. K. Elwood, N. Irges and P. Ramond, Phys. Rev. Lett. **81** (1998), 5064.
N. Irges, S. Lavignac and P. Ramond, Phys. Rev. **D58** (1998), 035003.
- [12] For example, W. Grimus, A. S. Joshipura, L. Lavoura and M. Tanimoto, Eur. Phys. J. **C36** (2004), 227.
- [13] S. Pakvasa and H. Sugawara, Phys. Lett. B **73** (1978), 61.
D. Wyler, Phys. Rev. D **19** (1979), 330.
L. J. Hall and H. Murayama, Phys. Rev. Lett. **75** (1995), 3985.
C. D. Carone, L. J. Hall and H. Murayama, Phys. Rev. D **53**, (1996) 6282.
R. Dermisek and S. Raby, Phys. Rev. D **62** (2000), 015007.
R. N. Mohapatra, A. Perez-Lorenzana and C. A. de Sousa Pires, Phys. Lett. B **474** (2000), 355.
J. Kubo, A. Mondragon, M. Mondragon and E. Rodriguez-Jauregui, Prog. Theor. Phys. **109** (2003), 795 [Errata; **114** (2005), 287].
T. Kobayashi, J. Kubo and H. Terao, Phys. Lett. B **568** (2003), 83.
P. F. Harrison and W. G. Scott, Phys. Lett. B **557** (2003), 76.
J. Kubo, H. Okada and F. Sakamaki, Phys. Rev. D **70** (2004), 036007.
S.-L. Chen, M. Frigerio and E. Ma, Phys. Rev. D **70** (2004), 073008 [Errata; D **70** (2004), 079905].
W. Grimus and L. Lavoura, J. High Energy Phys. **08** (2005), 013.
Y. Koide, Phys. Rev. D **73** (2006), 057901.

- [14] N. Haba and K. Yoshioka, Nucl. Phys. B **739** (2006), 254.
- [15] S. Pakvasa and H. Sugawara, Phys. Lett. B **82** (1979), 105.
 Y. Yamanaka, H. Sugawara and S. Pakvasa, Phys. Rev. D **25** (1982), 1895 [Errata; D **29** (1984), 2135].
 C. Hagedorn, M. Lindner and R. N. Mohapatra, J. High Energy Phys. **06** (2006), 042.
- [16] W. Grimus and L. Lavoura, Phys. Lett. B **572** (2003), 189.
 W. Grimus, A. S. Joshipura, S. Kaneko, L. Lavoura and M. Tanimoto, J. High Energy Phys. **07** (2004), 078.
 W. Grimus, A. S. Joshipura, S. Kaneko, L. Lavoura, H. Sawanaka and M. Tanimoto, Nucl. Phys. B **713** (2005), 151.
- [17] K. S. Babu and J. Kubo, Phys. Rev. D **71** (2005), 056006.
 Y. Kajiyama, E. Itou and J. Kubo, Nucl. Phys. B **743** (2006), 74.
- [18] M. Frigerio, S. Kaneko, E. Ma and M. Tanimoto, Phys. Rev. D **71** (2005), 011901.
- [19] E. Ma and G. Rajasekaran, Phys. Rev. D **64** (2001), 113012.
 E. Ma, Mod. Phys. Lett. A **17** (2002), 2361.
 K. S. Babu, E. Ma and J. W. F. Valle, Phys. Lett. B **552** (2003), 207.
 M. Hirsch, J. C. Romao, S. Skadhauge, J. W. F. Valle and A. Villanova del Moral, Phys. Rev. D **69** (2004), 093006.
 A. Zee, Phys. Lett. B **630** (2005), 58.
 G. Altarelli and F. Feruglio, Nucl. Phys. B **720** (2005), 64.
 X.-G. He, Y.-Y. Keum and R.R. Volkas, J. High Energy Phys. **04** (2006), 039.
 E. Ma, H. Sawanaka and M. Tanimoto, hep-ph/0606103.
- [20] E. Derman and H. S. Tsao, Phys. Rev. D **20** (1979), 1207.
 D. Chang, W. Y. Keung and G. Senjanovic, Phys. Rev. D **42** (1990), 1599.
 D. B. Kaplan and M. Schmaltz, Phys. Rev. D **49** (1994), 3741.
 P. H. Frampton and T. W. Kephart, Int. J. Mod. Phys. A **10** (1995), 4689.
 A. Aranda, C. D. Carone and R. F. Lebed, Phys. Lett. B **474** (2000), 170.
 N. Haba, A. Watanabe and K. Yoshioka, Phys. Rev. Lett. **97** (2006), 041601.
 C. Hagedorn, M. Lindner and F. Plentinger, Phys. Rev. D **74** (2006), 025007.
- [21] H. Georgi and C. Jarlskog, Phys. Lett. B **86** (1979), 297.
 S. Dimopoulos, L. J. Hall and S. Raby, Phys. Rev. Lett. **68** (1992), 1984.
- [22] K. S. Babu and Q. Shafi, Phys. Rev. D **47** (1993), 11.
 K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **74** (1995), 2418.
- [23] S. Eidelman et al. [Particle Data Group Collaboration], Phys. Lett. B **592** (2004), 1.
- [24] K. Anderson et al., hep-ex/0402041.
- [25] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, Nucl. Phys. B **477** (1996), 321.